

Circumcision Procedure Note

Date of Procedure:		
Pre-Op Diagnosis:	Parent(s) desire infant circumcision	
Post-Op Diagnosis:	Status post infant circumcision	
	Circumcision not done due to:	
Procedure:	Infant circumcision using Gomco clamp	
	Infant circumcision using Mogan clamp	
	Infant circumcision using Plastibell	
Anesthesia/Analgesia:	Ring block	
	Dorsal block	
	Penile block	
	Topical	
	cc of 1% lidocaine without epinephrine	
	Acetaminophen 12 mg/kg po 30 min prior to procedure	
Other:		
Surgeon:	Resident:	
	Attending:	
Estimated Blood Loss:	Minimal	
	Other:	
Indications for the Procedure:	Parent(s) desired newborn circumcision of their male infant. Prior to the procedure, the infant was examined and has no signs of hypospadias or illness.	
	The infant is term.	
	The infant was preterm, but has achieved an adequate weight.	
Risks, benefits and alternatives:	Were discussed with the parent(s) prior to the procedure, and informed consent was obtained. Signed consent form is in the infant's medical record. Discussion included, but was not limited to: no medical necessity for the procedure, possible bleeding, infection, damage to the penis or adjacent organs, possible poor cosmetic result and possible need for repeat procedure. All their questions were answered.	
	Parent(s) were informed that third party payers may not pay all or part of the circumcision fee. The parent(s) have agreed to be financially responsible for any unpaid portion of the circumcision fee.	
Complications:	None	
	Other:	
Procedure:	Area was prepped and draped in sterile fashion. Local anesthesia was administered as documented above under Anesthesia. After allowing sufficient time for the anesthesia to take effect, circumcision was performed in the usual sterile fashion using a	
	cm Gomco clamp	
	cm Plastibell	
	Mogan clamp	
	Good cosmesis and hemostasis was obtained. Vaseline gauze was applied. Infant tolerated the procedure well and was returned to the	
	Newborn Nursery	
	Mother's room	
in excellent condition. Mother was instructed how to care for the circumcision site.		